

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and protective (rubber or chemical resistant) gloves. Harmful if swallowed or if absorbed through the skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

TACTICAL STRIKE™ is a concentrated, one-step, Hospital Use, disinfectant that is effective against a broad spectrum of bacteria, is virucidal* including HIV-1, the AIDS Virus and Hepatitis B Virus and Hepatitis C Virus, fungicidal, and inhibits the growth of mold and mildew, and their odors, when used as directed. **TACTICAL STRIKE™** is a versatile disinfectant and sanitizer for hard, nonporous nonfood contact surfaces in Veterinarian, Animal care and Farm Premise applications. Kills HIV virus in 1 minute contact time. Kills flu virus in 1 minute contact time. Kills Germs, Bacteria, Viruses* on pre-cleaned, environmental surfaces. **Kills 99.9% SARS-Related Coronavirus* (SARS-CoV-2) (USA WA 1/2020) the virus that causes COVID-19 in 3 minutes.** Kills Norovirus in 5 minutes at 1: 64 dilution 2 oz. per gallon. **TACTICAL STRIKE™** is effective against Enterovirus D68 in 5 minutes at 1:64 dilution 2 oz. per gallon dilution kills Rotavirus in 3 minutes. Leaves a clean fresh scent. Sanitizes* 99.9% of bacteria in 15 seconds. *on hard, non-porous, non-food contact surfaces.

This product may be applied through low-pressure sprayers.

Use **TACTICAL STRIKE™** to clean and disinfect nonporous athletic mats, wrestling mats, gymnastic mats, exercise equipment and training tables, for non-scratch cleaning of showers and tubs, shower doors and curtains, fixtures and toilet bowls, for sanitizing and disinfecting of ultrasound transducers, probes, mammography compressor plates and other hard nonporous surfaces. Use **TACTICAL STRIKE™** to clean and disinfect non-porous personal protective safety equipment: protective headgear, hard hats, half-mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, hearing protectors and ear muffs. Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Precaution: Cleaning at 120°F temperature will avoid overheating and distortion of the personal safety equipment which would necessitate replacement). For Farm Premise Sanitation of hard nonporous surfaces: floors, walls, feed racks, mangers, troughs, automatic feeders, fountains and waterers, forks, shovels, scrapers and other nonporous surfaces in barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Homes, Veterinary Clinics, Kennels, Offices, Beauty Salons, Locker Rooms, Health Spas, Conference rooms, Elevators, Lounges, Day Care Centers, Theaters, Farms, Auditoriums and Churches.

Sanitizes fabric surfaces in 5 minutes. **TACTICAL STRIKE™** is a proven soft surface sanitizer against: Staphylococcus aureus, Enterobacter aerogenes. **TACTICAL STRIKE™** solution can be used 1 year after dilution when stored in a sealed container.

Use **TACTICAL STRIKE™** in: Bathroom(s), Restroom(s), Homes, Clinics, Dental offices, Health Care Facilities, Hospital(s) Medical Offices, Nursing homes, Sick Rooms, Day care centers, Nurseries, Convenience stores, Institutional kitchens, USDA inspected food processing facilities, Athletic facilities, Barber shops, Business and office buildings, Colleges, Correctional facilities, Hotels, Public facilities, Public rest rooms, Shower and bath areas, Animal laboratories, Farms, Zoos, Airplanes, Airports, Ambulances, Buses, Cars, Emergency (Police, EMS, Fire, Rescue) vehicles, Ships, Taxis, Trains, Transportation terminals. **Use TACTICAL STRIKE™ on soft surfaces of:** Upholstery, Fabric.

Use **TACTICAL STRIKE™** on washable hard, nonporous surfaces of: Bathroom fixtures, Countertops, Finished Floors, Tables, Toilet Exterior, Tiles, Tub's, glazed, Urinals, Walls, Non-critical medical device surfaces: Hospital Beds. Other hard nonporous surfaces made of: Glazed ceramic, Glazed enameled surfaces, Glazed porcelain, Laminated surfaces, Metal, Plastic, Sealed stone, Stainless steel, Upholstery, vinyl and plastic, Woodwork, finished.

TACTICAL STRIKE™ is a sanitizer for hard nonporous non-food contact surfaces at 1:128 dilution (1 oz. per gallon) tested according to the method for Sanitizers – Non-food Contact surfaces, modified in the presence of 250 ppm hard water plus 5% organic serum in 15 seconds against: Enterobacter aerogenes, Klebsiella pneumoniae, Listeria monocytogenes, Staphylococcus aureus, Staphylococcus aureus-Methicillin-Resistant (MRSA)

TACTICAL STRIKE™ is a sanitizer for soft, non-food contact surfaces at 1:128 dilution (1 oz. per gallon) according to the method for Sanitizers Recommended for Soft Non-food Contact surfaces, modified in the presence of 250 ppm hard water plus 5% organic serum in 5 minutes against: Enterobacter aerogenes, Staphylococcus aureus

Dilution:
Disinfection (1:128) 1 oz. per gallon of water
Disinfection (1:64) 2 oz. per gallon of water
Disinfection (1:32) 4 oz. per gallon of water
Virucidal: 1 - 2 oz. per gallon of water
Sanitizer (1:128) 1 oz. per gallon of water

 **SNEE CHEMICAL COMPANY**
5565 PEPSI STREET
HARAHAN, LOUISIANA

TACTICAL STRIKE™

Unscented Concentrated Disinfectant Cleaner Deodorant

This product contains no phosphates.

Staphylocidal • Pseudomonacidal • Bactericidal • Fungicidal • Virucidal*

TACTICAL STRIKE™ is effective against:

1 MINUTE CONTACT TIME:

Viruses:

*HIV-1 (AIDS Virus)

*Influenza Type A / Hong Kong Influenza

3 MINUTE CONTACT TIME:

Bacteria:

Klebsiella pneumoniae (Klebsiella)

Klebsiella pneumoniae, Carbapenem-resistant

Listeria monocytogenes

Proteus vulgaris

Serratia marcescens at 1:64 dilution (2 oz. per gallon of water)

Shigella dysenteriae (Shigella)

Shigella flexneri serotype 1B

Shigella sonnei

Staphylococcus aureus -

Community Associated Methicillin-Resistant (CA-MRSA) (NRS384) (USA300)

Staphylococcus aureus -

Community Associated Methicillin-Resistant (CA-MRSA) (NRS123) (USA400)

Staphylococcus aureus -

Methicillin-Resistant (MRSA)

Staphylococcus aureus - Multi-Drug Resistant

Staphylococcus aureus - Vancomycin Intermediate Resistant – (VISA)

Streptococcus pyogenes (Strep) (a cause of scarlet fever)

Yibrio cholerae

Veroina enterocolitica

Viruses:

*SARS-Related Coronavirus 2 (SARS-CoV-2) (cause of COVID-19)

*Hepatitis B Virus (HBV)

*Hepatitis C Virus (HCV)

*Herpes Simplex Virus Type 1

*Herpes Simplex Virus Type 2

*Human Coronavirus

*Respiratory Syncytial Virus (RSV)

*Rotavirus

*Vaccinia

Animal Viruses:

Canine Distemper Virus

Newcastle's Disease Virus

Active Ingredients:

Octyl decyl dimethyl ammonium chloride 3.34%

Dioctyl dimethyl ammonium chloride 1.33%

Didecyl dimethyl ammonium chloride 2.00%

Alkyl (C₁₄, 50%; C₁₂, 40%; C₁₆, 10%) dimethyl benzyl ammonium chloride 4.45%

Other Ingredients: 88.88%

Total: 100.00%

3 MINUTE CONTACT TIME:

Bacteria:

Pseudomonas aeruginosa (Pseudomonas)

Salmonella enterica (Salmonella)

Staphylococcus aureus (Staph)

Acinetobacter baumannii (Acinetobacter)

Bordetella bronchiseptica (Kennel cough)

Bordetella pertussis (whooping cough)

Campylobacter jejuni (Campylobacter)

Enterobacter aerogenes at 1:64 dilution 2 oz. per gallon of water)

Enterococcus faecalis (Enterococcus)

Enterococcus faecalis - Vancomycin resistant (VRE)

ESBL Escherichia coli (Extended spectrum beta-lactamase producing E. coli)

Escherichia coli (E. coli)

Escherichia coli O157:H7

5 MINUTE CONTACT TIME:

Viruses:

*Enterovirus D68 – at 1:64 dilution

(2 oz. per gallon of water)

*Norwalk Virus, Norovirus, Feline Calicivirus –

at 1:64 dilution (2 oz. per gallon of water)

Animal Virus:

Feline Calicivirus at 1:64 dilution

(2 oz. per gallon of water)

10 MINUTE CONTACT TIME:

Bacteria:

Legionella pneumophila

Viruses:

*Adenovirus Type 5 (Adenovirus) at 4 oz. per gallon

*Adenovirus Type 7 (Adenovirus) at 4 oz. per gallon

*SARS Associated Coronavirus (SARS)

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

Preparation of Disinfectant Use-Solution:

Add 1 oz. per gallon of water 1:128 For heavy duty use, add 4 oz. per gallon of water 1:32. Use 2 oz. per gallon of water to kill Norovirus, Enterobacter aerogenes, Serratia marcescens, and Enterovirus D68. Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7.

DISINFECTION / VIRUCIDAL* / FUNGICIDAL / MOLD AND MILDEW CONTROL DIRECTIONS:

Spray use-solution 6-8 inches from hard, nonporous surfaces, thoroughly wetting surfaces. Treated surfaces must remain visibly wet for 3 minutes. Do not breathe spray. For disinfection against HIV and Influenza type A/Hong Kong, use a 1 minute contact time. For disinfection against Enterovirus and Norovirus (Norwalk virus) use 5 minute contact time. For disinfection against (Legionella pneumophila), Adenovirus Type 5, Adenovirus Type 7 and SARS Associated Coronavirus use 10 minute contact time. Wipe dry with a cloth, sponge or mop or allow to air dry. For visibly soiled areas, a preliminary cleaning is required. To disinfect food processing facilities: Spray solution 6-8 inches from the surfaces and allow to remain wet for 3 minutes. Rinse food contact surfaces with potable water prior to reuse.

Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes.

Prepare a fresh solution daily or more often if the use solution becomes visibly diluted or soiled.

For fungicidal activity: **TACTICAL STRIKE™** is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. **TACTICAL STRIKE™** is a one-step fungicide at 1 oz. per gallon in 250 ppm hard water and 5% serum.

For mold and mildew: At 1 oz. per gallon, **TACTICAL STRIKE™** will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears.

This product is not for use on medical device surfaces.

TO SANITIZE SOFT SURFACES:

Add 1 oz. per gallon of water 1:128 or 1 - 2 oz. per gallon of water or 1 – 4 oz. per gallon of water. Apply use-solution with a sprayer. Hold bottle 6" – 8" from surface and spray until fabric is wet. Fabric must remain visibly wet for 5 minutes. Allow fabric to Air dry. Reapply as necessary. Visibly soiled areas must be cleaned prior to sanitizing.

***KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

Special Instructions for Cleaning and Decontamination against HIV-1 or HBV or HCV on Surfaces / Objects Soiled With Blood / Body Fluids.

Personal Protection: Cleanup must always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning Procedure: Blood and other body fluids containing HIV-1 or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of **TACTICAL STRIKE™**.

Contact Time: Leave surface wet for 1 minute for HIV-1 and 3 minutes for HBV and HCV (with a 1 – 4 oz.) per gallon use-solution. Use 2 oz. per gallon of water to kill Norovirus. Use contact time indicated on the label for disinfection against all other viruses, bacteria and fungi claimed.

Disposal of Infectious Material: Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

NON-ACID TOILET BOWL AND URINAL DISINFECTANT / CLEANER DIRECTIONS:

Add (1 – 4 oz.) product to toilet water. Use 2 oz. per gallon of water to kill Norovirus, Enterobacter aerogenes and Serratia marcescens. Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Brush or swab thoroughly, allow to stand for contact time indicated on the label and flush.

For Heavy Duty Use: Add 4 oz. product to toilet water. Brush or swab thoroughly, allow to stand for contact time indicated on the label and flush.

SALON / BARBER INSTRUMENTS AND TOOLS DISINFECTION (BACTERICIDE) / VIRUCIDE / FUNGICIDE DIRECTIONS:

Mix (1 – 4 oz.) of **TACTICAL STRIKE™** per gallon of water to disinfect hard, nonporous instruments and tools. Use 2 oz. per gallon of water to kill Norovirus, Enterobacter aerogenes and Serratia marcescens. Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Spray 6-8 inches away from the surface of combs, brushes, plastic rollers, manicure and other salon instruments and tools and allow them remain wet for contact time indicated on the label. For visibly soiled instruments and tools, a preliminary cleaning is required. Rinse thoroughly and dry before use.

Note: Stainless steel shears and instruments must be rinsed and dried and kept in a clean, non-contaminated receptacle. Prolonged contact with the disinfectant solution may cause damage to metal instruments.

VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOP / KENNELS /VIRUCIDAL* DISINFECTION DIRECTIONS:

For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces by spraying 6-8 inches away from the surface with a use-solution of (1 – 4 oz.) of **TACTICAL STRIKE™** per gallon of water for contact time indicated on the label. Use 2 oz. per gallon of water to kill Feline Calicivirus. Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Wipe or allow to air dry. Immerse all animal halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used for removing litter and manure. Rinse all surfaces that come in contact with food, including equipment used for feeding or watering with potable water before reuse. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

FARM PREMISE USE:

Remove all animals and feed from premises, vehicles, coops, crates and other enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces by spraying 6-8 inches away from the surface with a use-solution of (1 – 4 oz.) of **TACTICAL STRIKE™** per gallon of water for contact time indicated on the label. Use 2 oz. per gallon of water to kill Feline Calicivirus. Use 4 oz. per gallon of water to kill Adenovirus Type 5 and Type 7. Spray all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure. Ventilate treated buildings, vehicles, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

WATER DAMAGE RESTORATION AGAINST ODOR-CAUSING BACTERIA AND FUNGI FOR HOME, INSTITUTIONAL, AND INDUSTRIAL USE:

This product is particularly suitable for use in water damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushions, subfloors, drywall, trim and frame lumber, tackless strip and paneling. Using appropriate dilution, saturate the affected materials with enough product to remain wet for at least 3 minutes.

Sewer Backup and River Flooding: During mitigation procedures, dilute 1 – 2 oz. of the concentrate per gallon of water, allowing for the diluting effect of absorbed water within saturated materials. Remove [heavy soil] along with non-salvageable materials. Saturate all affected areas with a coarse spray device before and after cleaning and extraction.

Carpets, Carpet Cushions and Other Porous Materials such as Subfloors, Drywall, Trim and Frame Lumber, Tackless Strip and Paneling: For water damage from a clean source, extract excess water. Test hidden area for colorfastness. Dilute 1 – 2 oz. of the concentrate per gallon of water, allowing for the diluting effect of absorbed water within saturated materials. Remove excess filth or heavy soil. Saturate all affected areas with a coarse spray device. Roll brush or agitate into the materials and allow materials to remain wet for 3 minutes. Follow with a thorough extraction. Dry rapidly and thoroughly.

Special instructions For Cleaning Carpet Against Odor Causing Bacteria: Use this product in industrial and institutional areas such as motels, hotels, nursing homes, schools, and hospitals. Use on wet-cleanable synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for colorfastness. **For portable extraction units:** Mix 1 oz. of concentrate per gallon of water. **For truck mounted extraction machines:** Mix 1 – 2 oz. of concentrate per gallon of water and meter at 4 gallons per hour.

TO CLEAN AND DEODORIZE: Mix 1 oz. per gallon of water to clean and deodorize surfaces. Apply using a sprayer. Wipe or allow to air dry.

DEODORIZING DIRECTIONS:

Spray Solutions: 1 - 2 oz. per gallon of water eliminates undesirable odors. Allow surfaces to air dry.

Carpets: 1 - 2 oz. per gallon of water eliminates odors associated with urine, vomit, smoke, and mildew. Reapply using a spray device after cleaning for freshening effects. Allow to dry.

Sewage Backup, Water Damage: 2 – 4 oz. per gallon of water or desired concentration. Spray over affected areas before and after cleaning and extraction. Allow ten minutes contact time. Use proper ventilation; open windows.

Cleaning Solution: Dilute 1 - 2 oz. per gallon of water for cleaning of kitchen counters, tables, walls, bathrooms, toilet seats, floors and large routine applications. Apply to surfaces using trigger spray or other spray device. Remove excess liquid or allow surfaces to air dry.

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to children. Nonrefillable container. Do not reuse or refill this container. Wrap (container) and put in trash or offer for recycling if available.